

# Rectangular Parking Functions

Jean-Christophe Aval

Joint work with F. Bergeron (UQAM, Montréal)

IHP - 01/06/2017

# Main result

## Main result

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

## Main result

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

①  $\mathcal{P}_{ad,bd}$  ?

## Main result

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

- 1  $\mathcal{P}_{ad,bd}$  ?
- 2  $h_{bk}[ak \mathbf{x}]$  ?

## Main result

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

- 1  $\mathcal{P}_{ad,bd}$  ?
- 2  $h_{bk}[ak \mathbf{x}]$  ?
- 3 Frob ?

# Outline

- 1 Main result
- 2 Rectangular parking functions
- 3 Symmetric functions
- 4 Frobenius characteristic
- 5 Proof of the main result
- 6 Consequences
- 7 Generalization (Schröder)

# Outline

- 1 Main result
- 2 Rectangular parking functions**
- 3 Symmetric functions
- 4 Frobenius characteristic
- 5 Proof of the main result
- 6 Consequences
- 7 Generalization (Schröder)




# Dyck paths

# Dyck paths

Dyck paths  
(square  $n \times n$ )


# Dyck paths

Dyck paths  
(square  $n \times n$ )



## Dyck paths


Dyck paths  
(square  $n \times n$ )



Catalan numbers  
 $\frac{1}{n+1} \binom{2n}{n}$

## Dyck paths

Dyck paths  
(square  $n \times n$ )




Catalan numbers  
 $\frac{1}{n+1} \binom{2n}{n}$

Dyck paths  
(rational  $a \times b$ )  
( $a \wedge b = 1$ )


## Dyck paths

Dyck paths  
(square  $n \times n$ )




Catalan numbers  
 $\frac{1}{n+1} \binom{2n}{n}$

Dyck paths  
(rational  $a \times b$ )  
( $a \wedge b = 1$ )




## Dyck paths

Dyck paths  
(square  $n \times n$ )



Catalan numbers  
 $\frac{1}{n+1} \binom{2n}{n}$

Dyck paths  
(rational  $a \times b$ )  
( $a \wedge b = 1$ )



Rational  
Catalan numbers  
 $\frac{1}{a+b} \binom{a+b}{a}$

# Rectangular Dyck paths




# Rectangular Dyck paths


Dyck paths  
(in a rectangle  $m \times n$ )

# Rectangular Dyck paths

Dyck paths  
(in a rectangle  $m \times n$ )




# Rectangular Dyck paths



Dyck paths  
(in a rectangle  $m \times n$ )

$$(m, n) = (ad, bd), a \wedge b = 1$$

# Rectangular Dyck paths



Dyck paths  
(in a rectangle  $m \times n$ )

$$(m, n) = (ad, bd), a \wedge b = 1$$

$\mathcal{D}_{m,n}$  set of (rectangular) Dyck paths  $m \times n$ .


# Rectangular parking functions

# Rectangular parking functions


Rectangular parking functions  
( $m \times n$ )

# Rectangular parking functions

Rectangular parking functions  
( $m \times n$ )



# Rectangular parking functions




Rectangular parking functions  
( $m \times n$ )

$\mathcal{P}_{m,n}$  set of (rectangular) parking functions  $m \times n$



# Rectangular parking functions




Rectangular parking functions  
( $m \times n$ )

$\mathcal{P}_{m,n}$  set of (rectangular) parking functions  $m \times n$

$$|\mathcal{P}_{a,b}| = a^{b-1} \quad (a \wedge b = 1)$$

# Rectangular parking functions




Rectangular parking functions  
( $m \times n$ )

$\mathcal{P}_{m,n}$  set of (rectangular) parking functions  $m \times n$

$$|\mathcal{P}_{a,b}| = a^{b-1} \quad (a \wedge b = 1)$$

$$|\mathcal{P}_{n+1,n}| = (n+1)^{n-1}$$

# Rectangular parking functions




Rectangular parking functions  
( $m \times n$ )

$\mathcal{P}_{m,n}$  set of (rectangular) parking functions  $m \times n$

$$|\mathcal{P}_{a,b}| = a^{b-1} \quad (a \wedge b = 1)$$

$$|\mathcal{P}_{n+1,n}| = (n+1)^{n-1} \quad [\text{Konheim-Weiss 1966}]$$

# Rectangular parking functions



Rectangular parking functions  
( $m \times n$ )

$\mathcal{P}_{m,n}$  set of (rectangular) parking functions  $m \times n$

$\mathcal{L}_\alpha$  set of parking functions with (Dyck) path  $\alpha$

$$|\mathcal{P}_{a,b}| = a^{b-1} \quad (a \wedge b = 1)$$

$$|\mathcal{P}_{n+1,n}| = (n+1)^{n-1} \quad [\text{Konheim-Weiss 1966}]$$

# Outline

- 1 Main result
- 2 Rectangular parking functions
- 3 Symmetric functions**
- 4 Frobenius characteristic
- 5 Proof of the main result
- 6 Consequences
- 7 Generalization (Schröder)

# Symmetric functions

# Symmetric functions

$$\mathbf{x} = x_1, x_2, \dots$$

# Symmetric functions

$$\mathbf{x} = x_1, x_2, \dots$$

$$p_k(\mathbf{x}) = \sum_{i \geq 1} x_i^k$$



# Symmetric functions

$$\mathbf{x} = x_1, x_2, \dots$$

$$p_k(\mathbf{x}) = \sum_{i \geq 1} x_i^k$$

$$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$$

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}$$

# Symmetric functions

$$\mathbf{x} = x_1, x_2, \dots$$

$$p_k(\mathbf{x}) = \sum_{i \geq 1} x_i^k$$

$$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$$

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}$$

$$p_k(\mathbf{x} + \mathbf{y}) = p_k(\mathbf{x}) + p_k(\mathbf{y})$$

# Symmetric functions

$$\mathbf{x} = x_1, x_2, \dots$$

$$p_k(\mathbf{x}) = \sum_{i \geq 1} x_i^k$$

$$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$$

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}$$

$$p_k(\mathbf{x} + \mathbf{y}) = p_k(\mathbf{x}) + p_k(\mathbf{y})$$

$$p_k[2\mathbf{x}] = p_k(\mathbf{x} + \mathbf{x}) = 2 p_k(\mathbf{x})$$

## Symmetric functions

$$\mathbf{x} = x_1, x_2, \dots$$

$$p_k(\mathbf{x}) = \sum_{i \geq 1} x_i^k$$

$$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$$

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}$$

$$p_k(\mathbf{x} + \mathbf{y}) = p_k(\mathbf{x}) + p_k(\mathbf{y})$$

$$p_k[2\mathbf{x}] = p_k(\mathbf{x} + \mathbf{x}) = 2 p_k(\mathbf{x})$$

$$p_k[m\mathbf{x}] = m p_k(\mathbf{x})$$

# Complete symmetric functions

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$



# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots h_{\lambda_\ell}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots h_{\lambda_\ell}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$

$$h_2(\mathbf{x} + \mathbf{y}) =$$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots h_{\lambda_\ell}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$

$$h_2(\mathbf{x} + \mathbf{y}) = h_2(\mathbf{x}) + h_2(\mathbf{y}) + h_1(\mathbf{x}) h_1(\mathbf{y})$$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots h_{\lambda_\ell}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$

$$h_2(\mathbf{x} + \mathbf{y}) = h_2(\mathbf{x}) + h_2(\mathbf{y}) + h_1(\mathbf{x}) h_1(\mathbf{y})$$

$$h_2[2\mathbf{x}] =$$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots h_{\lambda_\ell}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$

$$h_2(\mathbf{x} + \mathbf{y}) = h_2(\mathbf{x}) + h_2(\mathbf{y}) + h_1(\mathbf{x}) h_1(\mathbf{y})$$

$$h_2[2\mathbf{x}] = h_2(\mathbf{x} + \mathbf{x}) = 2 h_2(\mathbf{x}) + h_{11}(\mathbf{x})$$

# Complete symmetric functions

$h_k$  complete symmetric function of degree  $k$

$$\sum_{k \geq 0} h_k t^k = \prod_{i \geq 1} \frac{1}{(1 - x_i t)}$$

$$h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots h_{\lambda_\ell}$$

$$h_2(\mathbf{x}) = x_1^2 + x_2^2 + \cdots + x_1 x_2 + x_1 x_3 + x_2 x_3 + \cdots$$

$$h_2(\mathbf{x} + \mathbf{y}) = h_2(\mathbf{x}) + h_2(\mathbf{y}) + h_1(\mathbf{x}) h_1(\mathbf{y})$$

$$h_2[2\mathbf{x}] = h_2(\mathbf{x} + \mathbf{x}) = 2 h_2(\mathbf{x}) + h_{11}(\mathbf{x})$$

$$h_2[m\mathbf{x}] = m h_2(\mathbf{x}) + \binom{m}{2} h_{11}(\mathbf{x})$$

# Complete symmetric functions

# Complete symmetric functions

$$h_n[m \mathbf{x}] =$$




## Complete symmetric functions

$$h_n[m \mathbf{x}] = \sum_{u_1 + u_2 + \cdots + u_m = n} h_{u_1}(\mathbf{x}) h_{u_2}(\mathbf{x}) \cdots h_{u_m}(\mathbf{x})$$

## Complete symmetric functions


$$h_n[m \mathbf{x}] = \sum_{u_1 + u_2 + \dots + u_m = n} h_{u_1}(\mathbf{x}) h_{u_2}(\mathbf{x}) \cdots h_{u_m}(\mathbf{x})$$



$(0, 0, 0, 2, 3, 0, 1, 0, 0)$

## Complete symmetric functions

$$h_n[m \mathbf{x}] = \sum_{u_1 + u_2 + \dots + u_m = n} h_{u_1}(\mathbf{x}) h_{u_2}(\mathbf{x}) \cdots h_{u_m}(\mathbf{x})$$




$(0, 0, 0, 2, 3, 0, 1, 0, 0)$

$\mathcal{B}_{m,n}$  set of rectangular paths  $m \times n$  (with last step = East)

## Complete symmetric functions

$$h_n[m \mathbf{x}] = \sum_{u_1 + u_2 + \dots + u_m = n} h_{u_1}(\mathbf{x}) h_{u_2}(\mathbf{x}) \cdots h_{u_m}(\mathbf{x})$$




$(0, 0, 0, 2, 3, 0, 1, 0, 0)$

$\mathcal{B}_{m,n}$  set of rectangular paths  $m \times n$  (with last step = East)

$$|\mathcal{B}_{m,n}| = \binom{m+n-1}{n}$$

## Complete symmetric functions

$$h_n[m \mathbf{x}] = \sum_{u_1 + u_2 + \dots + u_m = n} h_{u_1}(\mathbf{x}) h_{u_2}(\mathbf{x}) \cdots h_{u_m}(\mathbf{x})$$



 $\gamma \in \mathcal{B}_{m,n}$ 
 $(0, 0, 0, 2, 3, 0, 1, 0, 0)$ 
 $\rho(\gamma) = (3, 2, 1)$ 

$\mathcal{B}_{m,n}$  set of rectangular paths  $m \times n$  (with last step = East)

$$|\mathcal{B}_{m,n}| = \binom{m+n-1}{n}$$

## Complete symmetric functions

$$\begin{aligned}
 h_n[m \mathbf{x}] &= \sum_{u_1+u_2+\dots+u_m=n} h_{u_1}(\mathbf{x}) h_{u_2}(\mathbf{x}) \cdots h_{u_m}(\mathbf{x}) \\
 &= \sum_{\gamma \in \mathcal{B}_{m,n}} h_{\rho(\gamma)}(\mathbf{x}) \quad (h_0 = 1)
 \end{aligned}$$


 $(0, 0, 0, 2, 3, 0, 1, 0, 0)$ 
 $\gamma \in \mathcal{B}_{m,n}$ 
 $\rho(\gamma) = (3, 2, 1)$ 

$\mathcal{B}_{m,n}$  set of rectangular paths  $m \times n$  (with last step = East)

$$|\mathcal{B}_{m,n}| = \binom{m+n-1}{n}$$

# Outline

- 1 Main result
- 2 Rectangular parking functions
- 3 Symmetric functions
- 4 Frobenius characteristic**
- 5 Proof of the main result
- 6 Consequences
- 7 Generalization (Schröder)

# Symmetric group action



# Symmetric group action

$\mathcal{S}_n$  symmetric group

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$

$$\gamma \in \mathcal{B}_{m,n}$$




## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$




$$\gamma \in \mathcal{B}_{m,n}$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$



$$\gamma \in \mathcal{B}_{m,n}$$

$\mathcal{L}_\gamma$  increasing labellings of  $\gamma$




## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$



$$\gamma \in \mathcal{B}_{m,n}$$

$\mathcal{L}_\gamma$  increasing labellings of  $\gamma$


$$|\mathcal{L}_\gamma| =$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$



$$\gamma \in \mathcal{B}_{m,n}$$

$\mathcal{L}_\gamma$  increasing labellings of  $\gamma$


$$|\mathcal{L}_\gamma| = \binom{n}{\rho(\gamma)}$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
group of permutations over  $n$  letters

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$$

$$|\mathcal{S}_n| = n!$$



$$\gamma \in \mathcal{B}_{m,n}$$


$\mathcal{L}_\gamma$  increasing labellings of  $\gamma$

$$|\mathcal{L}_\gamma| = \binom{n}{\rho(\gamma)} = \frac{n!}{r_1! r_2! \cdots r_l!} \text{ for } \rho(\gamma) = (r_1, r_2, \dots, r_l)$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
 group of permutations over  $n$  letters  
 $\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$

$$|\mathcal{S}_n| = n!$$



$$\gamma \in \mathcal{B}_{m,n}$$

$\mathcal{L}_\gamma$  increasing labellings of  $\gamma$


$$|\mathcal{L}_\gamma| = \binom{n}{\rho(\gamma)} = \frac{n!}{r_1! r_2! \cdots r_l!} \text{ for } \rho(\gamma) = (r_1, r_2, \dots, r_l)$$

$$\sum_{\gamma \in \mathcal{B}_{m,n}} |\mathcal{L}_\gamma| = m^n$$

## Symmetric group action

$\mathcal{S}_n$  symmetric group  
 group of permutations over  $n$  letters  
 $\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$

$$|\mathcal{S}_n| = n!$$



$$\gamma \in \mathcal{B}_{m,n}$$

$$(6, 8, 4, 6, 6, 4)$$

$\mathcal{L}_\gamma$  increasing labellings of  $\gamma$

$$|\mathcal{L}_\gamma| = \binom{n}{\rho(\gamma)} = \frac{n!}{r_1! r_2! \cdots r_l!} \text{ for } \rho(\gamma) = (r_1, r_2, \dots, r_l)$$

$$\sum_{\gamma \in \mathcal{B}_{m,n}} |\mathcal{L}_\gamma| = m^n$$


# Symmetric group action

## Symmetric group action

$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels

# Symmetric group action


$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels





## Symmetric group action


$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels



$$\sigma = 425631$$

## Symmetric group action


$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels



$$\sigma = 425631$$

## Symmetric group action


$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels



$$\sigma = 425631$$

## Symmetric group action

$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels




$$\sigma = 425631$$

Transitive action on orbit =  $\mathcal{L}_\gamma$

## Symmetric group action

$S_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels




$$\sigma = 425631$$

Transitive action on orbit =  $\mathcal{L}_\gamma$

Stabilizer of  $\pi \in \mathcal{L}_\gamma$

## Symmetric group action

$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels




$$\sigma = 425631$$

Transitive action on orbit =  $\mathcal{L}_\gamma$

Stabilizer of  $\pi \in \mathcal{L}_\gamma = \mathcal{S}_2 \times \mathcal{S}_3 \times \mathcal{S}_1$

## Symmetric group action

$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels




$$\sigma = 425631$$

Transitive action on orbit =  $\mathcal{L}_\gamma$

$$\begin{aligned} \text{Stabilizer of } \pi \in \mathcal{L}_\gamma &= \mathcal{S}_2 \times \mathcal{S}_3 \times \mathcal{S}_1 \\ &= \mathcal{S}_{r_1} \times \mathcal{S}_{r_2} \times \cdots \times \mathcal{S}_{r_l} \text{ for } \rho(\gamma) = (r_1, r_2, \dots, r_l) \end{aligned}$$

## Symmetric group action

$\mathcal{S}_n$  acts on  $\mathcal{L}_\gamma$  by permuting the labels



$$\sigma = 425631$$

Transitive action on orbit =  $\mathcal{L}_\gamma$

$$\begin{aligned} \text{Stabilizer of } \pi \in \mathcal{L}_\gamma &= \mathcal{S}_2 \times \mathcal{S}_3 \times \mathcal{S}_1 \\ &= \mathcal{S}_{r_1} \times \mathcal{S}_{r_2} \times \cdots \times \mathcal{S}_{r_l} \text{ for } \rho(\gamma) = (r_1, r_2, \dots, r_l) \end{aligned}$$

→ Trivial action of  $\mathcal{S}_{r_1} \times \mathcal{S}_{r_2} \times \cdots \times \mathcal{S}_{r_l}$  induced on  $\mathcal{S}_n$



# Frobenius characteristic

# Frobenius characteristic

Action of  $\mathcal{S}_n$  with character  $\chi$

# Frobenius characteristic

Action of  $\mathcal{S}_n$  with character  $\chi$

Cyclic type

# Frobenius characteristic

Action of  $\mathcal{S}_n$  with character  $\chi$

Cyclic type  $\sigma = 425631$

# Frobenius characteristic

Action of  $\mathcal{S}_n$  with character  $\chi$

Cyclic type  $\sigma = 4\ 2\ 5\ 6\ 3\ 1 = (1, 4, 6)(2)(3, 5)$

# Frobenius characteristic

Action of  $\mathcal{S}_n$  with character  $\chi$

Cyclic type  $\sigma = 4\ 2\ 5\ 6\ 3\ 1 = (1, 4, 6)(2)(3, 5) \quad \lambda(\sigma) = (3, 2, 1)$

# Frobenius characteristic

Action of  $\mathcal{S}_n$  with character  $\chi$

Cyclic type  $\sigma = 4\ 2\ 5\ 6\ 3\ 1 = (1, 4, 6)(2)(3, 5) \quad \lambda(\sigma) = (3, 2, 1)$

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

# Frobenius characteristic



## Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

# Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism

# Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism
- $\text{Frob}(\chi^\lambda) = s_\lambda$  (Schur)

# Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism
- $\text{Frob}(\chi^\lambda) = s_\lambda$  (Schur)
- $\text{Frob}(\eta_n) = h_n$

# Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism
- $\text{Frob}(\chi^\lambda) = s_\lambda$  (Schur)
- $\text{Frob}(\eta_n) = h_n$     $\eta_\lambda = \text{Ind}_{\mathcal{S}_{\lambda_1} \times \dots \times \mathcal{S}_{\lambda_\ell}}^{\mathcal{S}_n} \eta_{\lambda_1} \times \dots \times \eta_{\lambda_\ell}$

## Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism
- $\text{Frob}(\chi^\lambda) = s_\lambda$  (Schur)
- $\text{Frob}(\eta_n) = h_n$     $\eta_\lambda = \text{Ind}_{\mathcal{S}_{\lambda_1} \times \dots \times \mathcal{S}_{\lambda_\ell}}^{\mathcal{S}_n} \eta_{\lambda_1} \times \dots \times \eta_{\lambda_\ell}$     $\text{Frob}(\eta_\lambda) = h_\lambda$

## Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism
- $\text{Frob}(\chi^\lambda) = s_\lambda$  (Schur)
- $\text{Frob}(\eta_n) = h_n$     $\eta_\lambda = \text{Ind}_{\mathcal{S}_{\lambda_1} \times \dots \times \mathcal{S}_{\lambda_\ell}}^{\mathcal{S}_n} \eta_{\lambda_1} \times \dots \times \eta_{\lambda_\ell}$     $\text{Frob}(\eta_\lambda) = h_\lambda$

Thus for  $\gamma \in \mathcal{B}_{m,n}$

$$\text{Frob}(\mathcal{L}_\gamma) = h_{\rho(\gamma)}$$

## Frobenius characteristic

$$\text{Frob}(\chi) = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n} \chi(\sigma) p_{\lambda(\sigma)}(\mathbf{x})$$

- Frob is an isomorphism
- $\text{Frob}(\chi^\lambda) = s_\lambda$  (Schur)
- $\text{Frob}(\eta_n) = h_n$     $\eta_\lambda = \text{Ind}_{\mathcal{S}_{\lambda_1} \times \dots \times \mathcal{S}_{\lambda_\ell}}^{\mathcal{S}_n} \eta_{\lambda_1} \times \dots \times \eta_{\lambda_\ell}$     $\text{Frob}(\eta_\lambda) = h_\lambda$

Thus for  $\gamma \in \mathcal{B}_{m,n}$

$$\text{Frob}(\mathcal{L}_\gamma) = h_{\rho(\gamma)}$$

$$\sum_{\gamma \in \mathcal{B}_{m,n}} \text{Frob}(\mathcal{L}_\gamma) = \sum_{\gamma \in \mathcal{B}_{m,n}} h_{\rho(\gamma)} = h_n[m \mathbf{x}]$$



# Outline

- 1 Main result
- 2 Rectangular parking functions
- 3 Symmetric functions
- 4 Frobenius characteristic
- 5 Proof of the main result**
- 6 Consequences
- 7 Generalization (Schröder)

# The formula

## The formula


$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$




$$\gamma \in \mathcal{B}_{m,n}$$



$$\gamma \in \mathcal{B}_{m,n}$$




$$\gamma \in \mathcal{B}_{m,n}$$






$$\gamma \in \mathcal{B}_{m,n}$$




$$\gamma \in \mathcal{B}_{m,n}$$




2 low points  $(1 \leq t \leq d)$

$$\gamma \in \mathcal{B}_{m,n}$$






2 low points  $(1 \leq t \leq d)$

Rotation


$\gamma \in \mathcal{B}_{m,n}$ 2 low points ( $1 \leq t \leq d$ )

Rotation




$\gamma \in \mathcal{B}_{m,n}$ 2 low points ( $1 \leq t \leq d$ )


Rotation





$\gamma \in \mathcal{B}_{m,n}$





2 low points  $(1 \leq t \leq d)$






Rotation




 $\gamma \in \mathcal{B}_{m,n}$ 

 2 low points  $(1 \leq t \leq d)$ 


Rotation







## Properties of the rotation

## Properties of the rotation

- for  $\gamma'$  image of  $\gamma$  by rotation :

## Properties of the rotation

- for  $\gamma'$  image of  $\gamma$  by rotation :

$$\rho(\gamma') = \rho(\gamma)$$

## Properties of the rotation

- for  $\gamma'$  image of  $\gamma$  by rotation :

$$\rho(\gamma') = \rho(\gamma) \implies \text{Frob}(\mathcal{L}_{\gamma'}) = \text{Frob}(\mathcal{L}_{\gamma})$$

## Properties of the rotation

- for  $\gamma'$  image of  $\gamma$  by rotation :

$$\rho(\gamma') = \rho(\gamma) \implies \text{Frob}(\mathcal{L}_{\gamma'}) = \text{Frob}(\mathcal{L}_{\gamma})$$

- the number  $t$  of low points is preserved

## Properties of the rotation

- for  $\gamma'$  image of  $\gamma$  by rotation :

$$\rho(\gamma') = \rho(\gamma) \implies \text{Frob}(\mathcal{L}_{\gamma'}) = \text{Frob}(\mathcal{L}_{\gamma})$$


- the number  $t$  of low points is preserved
- bijection :  $\mathcal{D}_{m,n}^t \times \llbracket 1, m \rrbracket \longrightarrow \mathcal{B}_{m,n}^t \times \llbracket 1, t \rrbracket$

## Properties of the rotation

- for  $\gamma'$  image of  $\gamma$  by rotation :

$$\rho(\gamma') = \rho(\gamma) \implies \text{Frob}(\mathcal{L}_{\gamma'}) = \text{Frob}(\mathcal{L}_{\gamma})$$

- the number  $t$  of low points is preserved
- bijection :  $\mathcal{D}_{m,n}^t \times \llbracket 1, m \rrbracket \longrightarrow \mathcal{B}_{m,n}^t \times \llbracket 1, t \rrbracket$



$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$



$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha) \quad a \text{ and } b \text{ fixed}$$

$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha)$$

$a$  and  $b$  fixed

$$\Phi_d^1(\mathbf{x}) = \sum_{\alpha \in \mathcal{D}_{ad,bd}} \text{Frob}(\mathcal{L}_\alpha)$$

primitive

$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha) \quad a \text{ and } b \text{ fixed}$$

$$\Phi_d^1(\mathbf{x}) = \sum_{\alpha \in \mathcal{D}_{ad,bd}} \text{Frob}(\mathcal{L}_\alpha)$$

primitive

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x})$$

$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha) \quad a \text{ and } b \text{ fixed}$$

$$\Phi_d^1(\mathbf{x}) = \sum_{\alpha \in \mathcal{D}_{ad,bd}} \text{Frob}(\mathcal{L}_\alpha)$$

primitive

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} \sum_{t>0} \sum_{\gamma \in \mathcal{B}_{m,n}^t} h_{\rho(\gamma)}$$

$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha) \quad a \text{ and } b \text{ fixed}$$

$$\Phi_d^1(\mathbf{x}) = \sum_{\alpha \in \mathcal{D}_{ad,bd}} \text{Frob}(\mathcal{L}_\alpha) \quad (m = ad)$$

primitive

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} \sum_{t>0} \sum_{\gamma \in \mathcal{B}_{m,n}^t} h_{\rho(\gamma)}$$

$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha) \quad a \text{ and } b \text{ fixed}$$

$$\Phi_d^1(\mathbf{x}) = \sum_{\alpha \in \mathcal{D}_{ad,bd}} \text{Frob}(\mathcal{L}_\alpha) \quad (m = ad)$$

primitive

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} \sum_{t>0} \sum_{\gamma \in \mathcal{B}_{m,n}^t} h_{\rho(\gamma)} = \frac{1}{ad} h_{bd}[ad \mathbf{x}]$$

$$\sum_{\alpha \in \mathcal{D}_{m,n}^t} \text{Frob}(\mathcal{L}_\alpha) \times m = \sum_{\gamma \in \mathcal{B}_{m,n}^t} \text{Frob}(\mathcal{L}_\gamma) \times t$$

$$\Phi_d^t(\mathbf{x}) := \sum_{\alpha \in \mathcal{D}_{ad,bd}^t} \text{Frob}(\mathcal{L}_\alpha) \quad a \text{ and } b \text{ fixed}$$

$$\Phi_d^1(\mathbf{x}) = \sum_{\alpha \in \mathcal{D}_{ad,bd}} \text{Frob}(\mathcal{L}_\alpha) \quad (m = ad)$$

primitive

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} \sum_{t>0} \sum_{\gamma \in \mathcal{B}_{m,n}^t} h_{\rho(\gamma)} = \frac{1}{ad} h_{bd}[ad \mathbf{x}]$$

$$\Phi_d^t(\mathbf{x}) = \sum_{\substack{c_1, c_2, \dots, c_t > 0 \\ c_1 + c_2 + \dots + c_t = d}} \Phi_{c_1}^1(\mathbf{x}) \Phi_{c_2}^1(\mathbf{x}) \cdots \Phi_{c_t}^1(\mathbf{x})$$

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} h_{bd}[ad \mathbf{x}]$$

$$\Phi_d^t(\mathbf{x}) = \sum_{\substack{c_1, c_2, \dots, c_t > 0 \\ c_1 + c_2 + \dots + c_t = d}} \Phi_{c_1}^1(\mathbf{x}) \Phi_{c_2}^1(\mathbf{x}) \cdots \Phi_{c_t}^1(\mathbf{x})$$



$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} h_{bd}[ad \mathbf{x}]$$

$$\Phi_d^t(\mathbf{x}) = \sum_{\substack{c_1, c_2, \dots, c_t > 0 \\ c_1 + c_2 + \dots + c_t = d}} \Phi_{c_1}^1(\mathbf{x}) \Phi_{c_2}^1(\mathbf{x}) \cdots \Phi_{c_t}^1(\mathbf{x})$$

$$P(\mathbf{x}; z) := \sum_{j>0} \Phi_j^1(\mathbf{x}) z^j$$

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} h_{bd}[ad \mathbf{x}]$$

$$\Phi_d^t(\mathbf{x}) = \sum_{\substack{c_1, c_2, \dots, c_t > 0 \\ c_1 + c_2 + \dots + c_t = d}} \Phi_{c_1}^1(\mathbf{x}) \Phi_{c_2}^1(\mathbf{x}) \cdots \Phi_{c_t}^1(\mathbf{x})$$

$$P(\mathbf{x}; z) := \sum_{j>0} \Phi_j^1(\mathbf{x}) z^j$$

$$\Phi_d^t(\mathbf{x}) = (P(\mathbf{x}; z))^t \Big|_{z^d}$$

$$\sum_{t>0} \frac{1}{t} \Phi_d^t(\mathbf{x}) = \frac{1}{ad} h_{bd}[ad \mathbf{x}]$$

$$\Phi_d^t(\mathbf{x}) = \sum_{\substack{c_1, c_2, \dots, c_t > 0 \\ c_1 + c_2 + \dots + c_t = d}} \Phi_{c_1}^1(\mathbf{x}) \Phi_{c_2}^1(\mathbf{x}) \cdots \Phi_{c_t}^1(\mathbf{x})$$

$$P(\mathbf{x}; z) := \sum_{j>0} \Phi_j^1(\mathbf{x}) z^j$$

$$\Phi_d^t(\mathbf{x}) = (P(\mathbf{x}; z))^t \Big|_{z^d}$$

$$\frac{1}{ad} h_{bd}[ad \mathbf{x}] = (-\log(1 - P(\mathbf{x}; z))) \Big|_{z^d}$$

$$P(\mathbf{x}; z) = 1 - \exp\left(-\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$P(\mathbf{x}; z) = 1 - \exp\left(-\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$\Phi_d^t(\mathbf{x}) = (P(\mathbf{x}; z))^t \Big|_{z^d}$$

$$P(\mathbf{x}; z) = 1 - \exp\left(-\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$\Phi_d^t(\mathbf{x}) = (P(\mathbf{x}; z))^t \Big|_{z^d} \implies \Phi_d(\mathbf{x}) = \sum_{t>0} \Phi_d^t(\mathbf{x}) = \left(\frac{1}{1 - P(\mathbf{x}; z)}\right) \Big|_{z^d}$$

$$P(\mathbf{x}; z) = 1 - \exp\left(-\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$\Phi_d^t(\mathbf{x}) = (P(\mathbf{x}; z))^t \Big|_{z^d} \implies \Phi_d(\mathbf{x}) = \sum_{t>0} \Phi_d^t(\mathbf{x}) = \left(\frac{1}{1 - P(\mathbf{x}; z)}\right) \Big|_{z^d}$$

$$\sum_{d \geq 0} \Phi_d(\mathbf{x}) z^d = \frac{1}{1 - P(\mathbf{x}; z)} = \exp\left(\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$P(\mathbf{x}; z) = 1 - \exp\left(-\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$\Phi_d^t(\mathbf{x}) = (P(\mathbf{x}; z))^t \Big|_{z^d} \implies \Phi_d(\mathbf{x}) = \sum_{t>0} \Phi_d^t(\mathbf{x}) = \left(\frac{1}{1 - P(\mathbf{x}; z)}\right) \Big|_{z^d}$$

$$\sum_{d \geq 0} \Phi_d(\mathbf{x}) z^d = \frac{1}{1 - P(\mathbf{x}; z)} = \exp\left(\sum_{k>0} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad, bd}) z^d = \exp\left(\sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k\right)$$



# Outline

- 1 Main result
- 2 Rectangular parking functions
- 3 Symmetric functions
- 4 Frobenius characteristic
- 5 Proof of the main result
- 6 Consequences**
- 7 Generalization (Schröder)

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad, bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\sum_{d \geq 0} |\mathcal{P}_{ad,bd}| z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} (ak)^{bk} z^k \right)$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\begin{aligned} \sum_{d \geq 0} |\mathcal{P}_{ad,bd}| z^d &= \exp \left( \sum_{k \geq 1} \frac{1}{ak} (ak)^{bk} z^k \right) \\ &= \exp \left( \sum_{k \geq 1} (ak)^{bk-1} z^k \right) \end{aligned}$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\begin{aligned} \sum_{d \geq 0} |\mathcal{P}_{ad,bd}| z^d &= \exp \left( \sum_{k \geq 1} \frac{1}{ak} (ak)^{bk} z^k \right) \\ &= \exp \left( \sum_{k \geq 1} (ak)^{bk-1} z^k \right) \end{aligned}$$

$$\sum_{d \geq 0} |\mathcal{D}_{ad,bd}| z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} \binom{ak + bk - 1}{bk} z^k \right)$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk} [ak \mathbf{x}] z^k \right)$$

$$\begin{aligned} \sum_{d \geq 0} |\mathcal{P}_{ad,bd}| z^d &= \exp \left( \sum_{k \geq 1} \frac{1}{ak} (ak)^{bk} z^k \right) \\ &= \exp \left( \sum_{k \geq 1} (ak)^{bk-1} z^k \right) \end{aligned}$$

$$\begin{aligned} \sum_{d \geq 0} |\mathcal{D}_{ad,bd}| z^d &= \exp \left( \sum_{k \geq 1} \frac{1}{ak} \binom{ak + bk - 1}{bk} z^k \right) \\ &= \exp \left( \sum_{k \geq 1} \frac{1}{ak + bk} \binom{ak + bk}{bk} z^k \right) \end{aligned}$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad, bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$



$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\text{Frob}(\mathcal{P}_{ad,bd}) = \sum_{\mu \vdash d} \frac{1}{m_\mu!} H_\mu^{a/b}(\mathbf{x})$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\text{Frob}(\mathcal{P}_{ad,bd}) = \sum_{\mu \vdash d} \frac{1}{m_\mu!} H_\mu^{a/b}(\mathbf{x})$$

$$\mu = (\mu_1, \mu_2, \dots, \mu_\ell) = (1^{m_1}, 2^{m_2}, \dots)$$

$$m_\mu! = \prod_j m_j!$$

$$H_\mu^{a/b}(\mathbf{x}) = \prod_{i=1}^{\ell} \frac{1}{a\mu_i} h_{a\mu_i}[b\mu_i \mathbf{x}]$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\text{Frob}(\mathcal{P}_{ad,bd}) = \sum_{\mu \vdash d} \frac{1}{m_\mu!} H_\mu^{a/b}(\mathbf{x})$$

$$\mu = (\mu_1, \mu_2, \dots, \mu_\ell) = (1^{m_1}, 2^{m_2}, \dots)$$

$$m_\mu! = \prod_j m_j!$$

$$H_\mu^{a/b}(\mathbf{x}) = \prod_{i=1}^{\ell} \frac{1}{a^{\mu_i}} h_{a\mu_i}[b\mu_i \mathbf{x}]$$

$$\text{Frob}(\mathcal{P}_{2a,2b}) = \frac{1}{2a} h_{2b}[2a \mathbf{x}] + \frac{1}{2} \left( \frac{1}{a} h_b[a \mathbf{x}] \right)^2$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\text{Frob}(\mathcal{P}_{ad,bd}) = \sum_{\mu \vdash d} \frac{1}{m_\mu!} H_\mu^{a/b}(\mathbf{x})$$

$$\mu = (\mu_1, \mu_2, \dots, \mu_\ell) = (1^{m_1}, 2^{m_2}, \dots)$$

$$m_\mu! = \prod_j m_j!$$

$$H_\mu^{a/b}(\mathbf{x}) = \prod_{i=1}^{\ell} \frac{1}{a\mu_i} h_{a\mu_i}[b\mu_i \mathbf{x}]$$

$$\text{Frob}(\mathcal{P}_{2a,2b}) = \frac{1}{2a} h_{2b}[2a \mathbf{x}] + \frac{1}{2} \left( \frac{1}{a} h_b[a \mathbf{x}] \right)^2$$

$$\text{Frob}(\mathcal{P}_{3a,3b}) = \frac{1}{3a} h_{3b}[3a \mathbf{x}] + \left( \frac{1}{2a} h_{2b}[2a \mathbf{x}] \right) \left( \frac{1}{a} h_b[a \mathbf{x}] \right) + \frac{1}{6} \left( \frac{1}{a} h_b[a \mathbf{x}] \right)^3$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\text{Frob}(\mathcal{P}_{ad,bd}) = \sum_{\mu \vdash d} \frac{1}{m_\mu!} H_\mu^{a/b}(\mathbf{x})$$

$$\mu = (\mu_1, \mu_2, \dots, \mu_\ell) = (1^{m_1}, 2^{m_2}, \dots)$$

$$m_\mu! = \prod_j m_j!$$

$$H_\mu^{a/b}(\mathbf{x}) = \prod_{i=1}^{\ell} \frac{1}{a\mu_i} h_{a\mu_i}[b\mu_i \mathbf{x}]$$

$$|\mathcal{D}_{2a,2b}| = \left( \frac{1}{2a+2b} \binom{2a+2b}{2a} \right) + \frac{1}{2} \left( \frac{1}{a+b} \binom{a+b}{a} \right)^2$$

$$|\mathcal{D}_{3a,3b}| = \left( \frac{1}{3a+3b} \binom{3a+3b}{3a} \right) + \left( \frac{1}{a+b} \binom{a+b}{a} \right) \left( \frac{1}{2a+2b} \binom{2a+2b}{2a} \right) + \frac{1}{6} \left( \frac{1}{a+b} \binom{a+b}{a} \right)^3$$

$$\sum_{d \geq 0} \text{Frob}(\mathcal{P}_{ad,bd}) z^d = \exp \left( \sum_{k \geq 1} \frac{1}{ak} h_{bk}[ak \mathbf{x}] z^k \right)$$

$$\text{Frob}(\mathcal{P}_{ad,bd}) = \sum_{\mu \vdash d} \frac{1}{m_\mu!} H_\mu^{a/b}(\mathbf{x})$$

$$\mu = (\mu_1, \mu_2, \dots, \mu_\ell) = (1^{m_1}, 2^{m_2}, \dots)$$

$$m_\mu! = \prod_j m_j!$$

$$H_\mu^{a/b}(\mathbf{x}) = \prod_{i=1}^{\ell} \frac{1}{a\mu_i} h_{a\mu_i}[b\mu_i \mathbf{x}]$$

$$|\mathcal{D}_{2a,2b}| = \left( \frac{1}{2a+2b} \binom{2a+2b}{2a} \right) + \frac{1}{2} \left( \frac{1}{a+b} \binom{a+b}{a} \right)^2$$

$$|\mathcal{D}_{3a,3b}| = \left( \frac{1}{3a+3b} \binom{3a+3b}{3a} \right) + \left( \frac{1}{a+b} \binom{a+b}{a} \right) \left( \frac{1}{2a+2b} \binom{2a+2b}{2a} \right) + \frac{1}{6} \left( \frac{1}{a+b} \binom{a+b}{a} \right)^3$$

[Bizley 1954]

# Outline

- 1 Main result
- 2 Rectangular parking functions
- 3 Symmetric functions
- 4 Frobenius characteristic
- 5 Proof of the main result
- 6 Consequences
- 7 Generalization (Schröder)**

# Generalization - Schröder parking functions




# Generalization - Schröder parking functions

Rectangular Schröder parking functions  
( $m \times n$ )


# Generalization - Schröder parking functions

Rectangular Schröder parking functions  
( $m \times n$ )



# Generalization - Schröder parking functions


Rectangular Schröder parking functions  
 $(m \times n)$



$\mathcal{S}_{m,n}^{(k)}$  set of (rectangular) Schröder parking functions  $m \times n$   
 with  $k$  diagonal steps

# Generalization - Schröder parking functions


Rectangular Schröder parking functions  
( $m \times n$ )



$\mathcal{S}_{m,n}^{(k)}$  set of (rectangular) Schröder parking functions  $m \times n$   
with  $k$  diagonal steps

$$\sum_{k \geq 0} \text{Frob } \mathcal{S}_{m,n}^{(k)}(\mathbf{x}) y^k$$

# Generalization - Schröder parking functions





Rectangular Schröder parking functions  
( $m \times n$ )

$\mathcal{S}_{m,n}^{(k)}$  set of (rectangular) Schröder parking functions  $m \times n$   
with  $k$  diagonal steps

$$\sum_{k \geq 0} \text{Frob } \mathcal{S}_{m,n}^{(k)}(\mathbf{x}) y^k = \text{Frob } \mathcal{P}_{m,n}(\mathbf{x} + y)$$

# Références

-  J.-C. Aval, F. Bergeron, *Interlaced Rectangular Parking Functions*, preprint (2015) arXiv:1503.03991.
-  J.-C. Aval, F. Bergeron, *Rectangular Schröder Parking Functions Combinatorics*, preprint (2016) arXiv:1603.09487.