

Graphon limit of random cographs

Mathilde Bouvel
(Loria, CNRS, Univ. Lorraine)

talk based on joint work with
Frédérique Bassino, Valentin Féray, Lucas Gerin,
Mickaël Maazoun and Adeline Pierrot

Arxiv:1907.08517


Séminaire de combinatoire Philippe Flajolet
April 1st, 2021



**We have to start somewhere:
Setting the problem**


Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.



Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:



Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:



Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:



Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:



Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:



Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:




Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:




Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:




Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:




Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:




Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.




Example:




Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.



Example:



Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.

Other characterizations: Cographs are

- the graphs **avoiding** $P_4 = \bullet\text{---}\bullet\text{---}\bullet\text{---}\bullet$ as an induced subgraph;
- the graphs whose modular decomposition **does not involve any prime graph**;
- the inversion graphs of **separable permutations**.

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.

Other characterizations: Cographs are

- the graphs **avoiding** $P_4 = \bullet\text{---}\bullet\text{---}\bullet\text{---}\bullet$ as an induced subgraph;
- the graphs whose modular decomposition **does not involve any prime graph**;
- the inversion graphs of **separable permutations**.

Prop.: **Cographs** form a **hereditary** class, *i.e.*, every induced subgraph of a cograph is a cograph.

Cographs

Definition: A **cograph** of size n is a graph $G = (V, E)$ with $|V| = n$ which can be constructed from graphs with one vertex by taking **disjoint unions** and **joins**.

Other characterizations: Cographs are

- the graphs **avoiding** $P_4 = \bullet\text{---}\bullet\text{---}\bullet$ as an induced subgraph;
- the graphs whose modular decomposition **does not involve any prime graph**;
- the inversion graphs of **separable permutations**.

Prop.: **Cographs** form a **hereditary** class, *i.e.*, every induced subgraph of a cograph is a cograph.

Central question for this talk:

What does a **uniform random cograph** of size n look like, when n goes to infinity?

Cographs: one of many cases

One question studied in the (huge!) random graph literature is the following:

For \mathcal{F} a family of graphs,
what is the typical behavior of a large graph in \mathcal{F} ?

Already studied for:

- perfect graphs [McDiarmid-Yolov, 2019]
- planar graphs [Noy, 2014]
- graphs embeddable in a surface of given genus [Dowden-Kang-Sprüssel, 2017]
- graphs in subcritical classes [Panagiotou-Stufler-Weller, 2016]
- large hereditary classes [Hatami-Janson-Szegedy, 2018]
- addable classes [McDiarmid-Steger-Welsh, 2006 ; Chapuy-Perarnau, 2019]

Which model?

Which discrete objects? Graphs may be

- **labeled**: in this case, vertices are numbered from 1 to n ;
- or **unlabeled**: vertices are indistinguishable.

Unlabeled graphs are equivalence classes of labeled graphs under the action of relabeling the vertices.

Here, we consider both **labeled and unlabeled cographs**.

Which model?

Which discrete objects? Graphs may be

- **labeled**: in this case, vertices are numbered from 1 to n ;
- or **unlabeled**: vertices are indistinguishable.

Unlabeled graphs are equivalence classes of labeled graphs under the action of relabeling the vertices.

Here, we consider both **labeled and unlabeled cographs**.



The **unlabeled version of a uniform labeled** cograph is **not** a **uniform unlabeled** cograph! Indeed, the number of distinct labelings depends on the internal symmetries of the graph.

Which model?

Which discrete objects? Graphs may be

- **labeled**: in this case, vertices are numbered from 1 to n ;
- or **unlabeled**: vertices are indistinguishable.

Unlabeled graphs are equivalence classes of labeled graphs under the action of relabeling the vertices.

Here, we consider both **labeled and unlabeled cographs**.



The **unlabeled version of a uniform labeled** cograph is **not** a **uniform unlabeled** cograph! Indeed, the number of distinct labelings depends on the internal symmetries of the graph.


Which continuous limit? We describe the limit in the space of **graphons**. Graphons were introduced by Lovász and co-authors in 2008, and attracted a lot of interest.

Graphons are appropriate to describe **limits of dense graphs**.

Some basics on graphons

What is (informally) a graphon?


In the discrete setting:



The **graphon** W_G associated with G is the equivalence class of w_G under the action of permuting rows and columns of M_G .

What is (informally) a graphon?

In the discrete setting:




The **graphon** W_G associated with G is the equivalence class of w_G under the action of permuting rows and columns of M_G .

Remarks:

- W_G does not depend on the order of the vertices chosen to write M_G .
- If G is **labeled**, W_G is the graphon of the unlabeled version of G .

What is (informally) a graphon?

In the discrete setting:



The **graphon** W_G associated with G is the equivalence class of w_G under the action of permuting rows and columns of M_G .

Continuous extension:

In general, a graphon is obtained **as above**, from a symmetric matrix M , possibly with **a continuum of rows and columns**, and with values in $[0, 1]$.

It is an equivalence class of symmetric functions from $[0, 1]^2 \rightarrow [0, 1]$ under the action of permuting rows and columns of M .

Characterization of (deterministic) graphon convergence

(Non-)definition:

The space of graphons is (up to technicalities) **metric**, for the **cut-distance** (and in addition is compact).

So, it makes sense to study **convergence** of a sequence of graphons $(W_n)_{n \geq 0}$ to a graphon W (for this cut-distance). We write $W_n \rightarrow W$.

Characterization of (deterministic) graphon convergence

(Non-)definition:

The space of graphons is (up to technicalities) **metric**, for the **cut-distance** (and in addition is compact).

So, it makes sense to study **convergence** of a sequence of graphons $(W_n)_{n \geq 0}$ to a graphon W (for this cut-distance). We write $W_n \rightarrow W$.

Typically, $W_n = W_{G_n}$, the graphon associated to a graph G_n , with the sequence of graphs (G_n) such that the **size of G_n grows to infinity** with n . In this case, we also write $G_n \rightarrow W$.

Characterization of (deterministic) graphon convergence

(Non-)definition:

The space of graphons is (up to technicalities) **metric**, for the **cut-distance** (and in addition is compact).

So, it makes sense to study **convergence** of a sequence of graphons $(W_n)_{n \geq 0}$ to a graphon W (for this cut-distance). We write $W_n \rightarrow W$.

Typically, $W_n = W_{G_n}$, the graphon associated to a graph G_n , with the sequence of graphs (G_n) such that the **size of G_n grows to infinity** with n . In this case, we also write $G_n \rightarrow W$.

Combinatorial characterization of convergence:

For (W_n) a sequence of graphons and W a graphon, $W_n \rightarrow W$ iff for any (finite) graph g , $\text{Dens}(g, W_n) \rightarrow \text{Dens}(g, W)$.

Let us now define the **density** of a graph g in a graphon.

Subgraph densities in graphs and graphons

Induced subgraph: The **subgraph** of $G = (V, E)$ induced by $V' \subset V$ is the graph with vertex set V' and edge set $E \cap (V' \times V')$.

Subgraph densities in graphs and graphons

Induced subgraph: The **subgraph** of $G = (V, E)$ induced by $V' \subset V$ is the graph with vertex set V' and edge set $E \cap (V' \times V')$.

Densities: Fix g a graph with k vertices, unlabeled.

- For a graph G , $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$, where $\text{SubGraph}_k(G)$ is the (random) subgraph of G induced by a k -tuple of i.i.d. uniform random vertices of G .

Subgraph densities in graphs and graphons

Induced subgraph: The **subgraph** of $G = (V, E)$ induced by $V' \subset V$ is the graph with vertex set V' and edge set $E \cap (V' \times V')$.

Densities: Fix g a graph with k vertices, unlabeled.

- For a graph G , $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$, where $\text{SubGraph}_k(G)$ is the (random) subgraph of G induced by a k -tuple of i.i.d. uniform random vertices of G .
- For a graphon W , $\text{Dens}(g, W) = \mathbb{P}(\text{Sample}_k(W) = g)$, where $\text{Sample}_k(W)$ is the (random) graph with k vertices v_1, \dots, v_k such that v_i and v_j are connected with probability $w(x_i, x_j)$, for x_1, \dots, x_k i.i.d. uniform random variables in $[0, 1]$ and $w : [0, 1]^2 \rightarrow [0, 1]$ a representative of W .

Subgraph densities in graphs and graphons

Induced subgraph: The **subgraph** of $G = (V, E)$ induced by $V' \subset V$ is the graph with vertex set V' and edge set $E \cap (V' \times V')$.

Densities: Fix g a graph with k vertices, unlabeled.

- For a graph G , $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$, where $\text{SubGraph}_k(G)$ is the (random) subgraph of G induced by a k -tuple of i.i.d. uniform random vertices of G .
- For a graphon W , $\text{Dens}(g, W) = \mathbb{P}(\text{Sample}_k(W) = g)$, where $\text{Sample}_k(W)$ is the (random) graph with k vertices v_1, \dots, v_k such that v_i and v_j are connected with probability $w(x_i, x_j)$, for x_1, \dots, x_k i.i.d. uniform random variables in $[0, 1]$ and $w : [0, 1]^2 \rightarrow [0, 1]$ a representative of W .

Remark: For any graph G , $\text{Dens}(g, W_G) = \text{Dens}(g, G)$.

Characterization of graphon convergence: the random case

Reminder: $G_n \rightarrow W$ iff $\text{Dens}(g, G_n) \rightarrow \text{Dens}(g, W)$ for all g , for (G_n) a sequence of (deterministic) graphs and W a (deterministic) graphon.

Characterization of graphon convergence: the random case

Reminder: $G_n \rightarrow W$ iff $\text{Dens}(g, G_n) \rightarrow \text{Dens}(g, W)$ for all g , for (G_n) a sequence of (deterministic) graphs and W a (deterministic) graphon.

What if we take (\mathbf{G}_n) random? ($\text{Dens}(g, \mathbf{G}_n)$ being then a real r.v.)

Characterization of graphon convergence: the random case

Reminder: $G_n \rightarrow W$ iff $\text{Dens}(g, G_n) \rightarrow \text{Dens}(g, W)$ for all g , for (G_n) a sequence of (deterministic) graphs and W a (deterministic) graphon.

What if we take (\mathbf{G}_n) random? ($\text{Dens}(g, \mathbf{G}_n)$ being then a real r.v.)

Theorem [Diaconis-Janson, 2008]:

The distribution of a random graphon \mathbf{W} is characterized by all expected subgraph densities $\mathbb{E}[\text{Dens}(g, \mathbf{W})]$ (for all g).

Theorem [Diaconis-Janson, 2008]:

Let (\mathbf{G}_n) be a sequence of random graphs. TFAE:

- \mathbf{G}_n tends in distribution to some random graphon, \mathbf{W} .
- For all g , $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$ converges to some value $\Delta_g \in [0, 1]$.

If this holds, in addition we have:

for all g , $\mathbb{E}[\text{Dens}(g, \mathbf{W})] = \Delta_g$, so that $(\Delta_g)_g$ characterizes \mathbf{W} .

Our work in a nutshell

Main result and proof strategy

Theorem:

For all n , let \mathbf{G}_n (resp. \mathbf{G}_n^u) be a uniform random labeled (resp. unlabeled) cograph with n vertices.

We have that \mathbf{G}_n (resp. \mathbf{G}_n^u) converges in distribution to a random graphon $\mathbf{W}^{1/2}$ called the **Brownian cographon** of parameter $1/2$.

Main result and proof strategy

Theorem:

For all n , let \mathbf{G}_n (resp. \mathbf{G}_n^u) be a uniform random labeled (resp. unlabeled) cograph with n vertices.

We have that \mathbf{G}_n (resp. \mathbf{G}_n^u) converges in distribution to a random graphon $\mathbf{W}^{1/2}$ called the **Brownian cographon** of parameter $1/2$.

Proof strategy (labeled case):

- Compute $\Delta_g = \mathbb{E}[\text{Dens}(g, \mathbf{W}^{1/2})]$ for all cographs g
- Express $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$ as a quotient of **coefficients of generating functions**, starting from

$$\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)] = \frac{\left| \left\{ (G, I) : \begin{array}{l} G=(V,E) \text{ labeled cograph of size } n, \\ I \in V^k \text{ which induces } g \end{array} \right\} \right|}{\left| \{ G \text{ labeled cograph of size } n \} \right| \cdot n^k}$$

- Estimate numerator and denominator using **analytic combinatorics**, in order to prove **convergence to Δ_g**

Main result and proof strategy

Theorem:

For all n , let \mathbf{G}_n (resp. \mathbf{G}_n^u) be a uniform random labeled (resp. unlabeled) cograph with n vertices.

We have that \mathbf{G}_n (resp. \mathbf{G}_n^u) converges in distribution to a random graphon $\mathbf{W}^{1/2}$ called the **Brownian cographon** of parameter $1/2$.

Proof strategy (labeled case):

- Compute $\Delta_g = \mathbb{E}[\text{Dens}(g, \mathbf{W}^{1/2})]$ for all cographs g
- Express $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$ as a quotient of **coefficients of generating functions**, starting from

$$\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)] = \frac{\left| \left\{ (G, I) : \begin{array}{l} G=(V,E) \text{ labeled cograph of size } n, \\ I \in V^k \text{ which induces } g \end{array} \right\} \right|}{\left| \{ G \text{ labeled cograph of size } n \} \right| \cdot n^k}$$

- Estimate numerator and denominator using **analytic combinatorics**, in order to prove **convergence to Δ_g**

Essential tool: encoding of cographs by **cotrees**.

Outline of (the rest of) the talk

About the main theorem:

- Cographs and [cotrees](#)
- Combinatorial proof of [convergence](#) in the [labeled](#) case
- Description of the [Brownian cographon](#)
- Corollary: [average degree distribution](#) in cographs
- How to deal with the [unlabeled](#) case

Additional results, questions, comments:

- [Vertex connectivity](#) distinguishes between the labeled and the unlabeled settings
- A parallel with [permutations](#), yielding new problems to work on
- [Independence number](#) of cographs



Cotrees and how to use them to compute

$$\lim_{n \rightarrow \infty} \mathbb{E}[\text{Dens}(g, G_n)]$$

Cographs and cotrees

A **labeled cotree** of size n is a rooted tree t with leaves $\{1, \dots, n\}$ s.t.

- t is not plane (*i.e.* the children of every internal node are not ordered);
- every internal node has at least two children;
- every internal node carries a decoration 0 or 1.





Mapping
from cotrees to cographs:
0 indicates disjoint union and
1 indicates join.

Cographs and cotrees

A **labeled cotree** of size n is a rooted tree t with leaves $\{1, \dots, n\}$ s.t.

- t is not plane (*i.e.* the children of every internal node are not ordered);
- every internal node has at least two children;
- every internal node carries a decoration 0 or 1.





Mapping
from cotrees to cographs:
0 indicates disjoint union and
1 indicates join.

Prop.: Vertices i and j are **connected** iff the **first common ancestor** of leaves i and j carries a 1.

Cographs and cotrees

A **labeled cotree** of size n is a rooted tree t with leaves $\{1, \dots, n\}$ s.t.

- t is not plane (*i.e.* the children of every internal node are not ordered);
- every internal node has at least two children;
- every internal node carries a decoration 0 or 1.



(Not one-to-one) Mapping
from cotrees to cographs:
0 indicates disjoint union and
1 indicates join.



Prop.: Vertices i and j are **connected** iff the **first common ancestor** of leaves i and j carries a 1.

Cographs and cotrees

A **labeled cotree** of size n is a rooted tree t with leaves $\{1, \dots, n\}$ s.t.

- t is not plane (*i.e.* the children of every internal node are not ordered);
- every internal node has at least two children;
- every internal node carries a decoration 0 or 1.

t is **canonical** if 0 and 1 alternate on every branch from the root to a leaf.



(Not one-to-one) Mapping
from cotrees to cographs:
0 indicates disjoint union and
1 indicates join.


Prop.: Vertices i and j are **connected** iff the **first common ancestor** of leaves i and j carries a 1.

Prop.: This mapping restricted to canonical cotrees is a **bijection**.

Induced subgraphs in cographs on their cotrees

t a canonical cotree \leftrightarrow G the corresponding cograph
a k -tuple $\ell = (\ell_1, \dots, \ell_k)$ of leaves \leftrightarrow a k -tuple I of vertices


Subtree of t induced by (ℓ_1, \dots, ℓ_k) = the cotree **labeled from** ℓ whose leaves are (ℓ_1, \dots, ℓ_k) and whose internal structure is inherited from t .



Induced subgraphs in cographs on their cotrees

t a canonical cotree \leftrightarrow G the corresponding cograph
a k -tuple $\ell = (\ell_1, \dots, \ell_k)$ of leaves \leftrightarrow a k -tuple I of vertices

Subtree of t induced by (ℓ_1, \dots, ℓ_k) = the cotree **labeled from** ℓ whose leaves are (ℓ_1, \dots, ℓ_k) and whose internal structure is inherited from t .



Prop.: Forgetting the labelings, the **subgraph of G induced by I** is the cograph corresponding to the **subtree of t induced by (ℓ_1, \dots, ℓ_k)**

Variations on $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$

- **Reminder:** $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$,
where $\text{SubGraph}_k(G)$ is the (random) subgraph of G
induced by a k -tuple of i.i.d. uniform random vertices of G .

Variations on $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$

- **Reminder:** $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$,
where $\text{SubGraph}_k(G)$ is the (random) subgraph of G
induced by a k -tuple of i.i.d. uniform random vertices of G .
- **Variant:** $\text{Dens}^{inj}(g, G) = \mathbb{P}(\text{SubGraph}_k^{inj}(G) = g)$,
where $\text{SubGraph}_k^{inj}(G)$ is the (random) subgraph of G
induced by a uniform random k -tuple of distinct vertices of G .

Variations on $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$

- **Reminder:** $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$,
where $\text{SubGraph}_k(G)$ is the (random) subgraph of G
induced by a k -tuple of i.i.d. uniform random vertices of G .
- **Variant:** $\text{Dens}^{inj}(g, G) = \mathbb{P}(\text{SubGraph}_k^{inj}(G) = g)$,
where $\text{SubGraph}_k^{inj}(G)$ is the (random) subgraph of G
induced by a uniform random k -tuple of distinct vertices of G .

Fact: $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)] \rightarrow \Delta_g$ iff $\mathbb{E}[\text{Dens}^{inj}(g, \mathbf{G}_n)] \rightarrow \Delta_g$.

Variations on $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$

- **Reminder:** $\text{Dens}(g, G) = \mathbb{P}(\text{SubGraph}_k(G) = g)$, where $\text{SubGraph}_k(G)$ is the (random) subgraph of G induced by a k -tuple of i.i.d. uniform random vertices of G .
- **Variante:** $\text{Dens}^{\text{inj}}(g, G) = \mathbb{P}(\text{SubGraph}_k^{\text{inj}}(G) = g)$, where $\text{SubGraph}_k^{\text{inj}}(G)$ is the (random) subgraph of G induced by a uniform random k -tuple of distinct vertices of G .

Fact: $\mathbb{E}[\text{Dens}(g, \mathbf{G}_n)] \rightarrow \Delta_g$ iff $\mathbb{E}[\text{Dens}^{\text{inj}}(g, \mathbf{G}_n)] \rightarrow \Delta_g$.

Notation: for all n , and all $k \leq n$,

$\mathbf{t}^{(n)}$ is a uniform random labeled canonical cotree of size n , and $\mathbf{t}_k^{(n)}$ is the subtree of $\mathbf{t}^{(n)}$ induced by a uniform k -tuple of distinct leaves.

For any cograph g , we have:

$$\mathbb{E}[\text{Dens}^{\text{inj}}(g, \mathbf{G}_n)] = \mathbb{P}(\text{SubGraph}_k^{\text{inj}}(\mathbf{G}_n) = g) = \sum \mathbb{P}(\mathbf{t}_k^{(n)} = t_0),$$
 where the sum runs over all cotrees t_0 corresponding to g .

Combinatorics of the labeled case:

Finding $\lim_{n \rightarrow \infty} \mathbb{P}(t_k^{(n)} = t_0)$

Notation:

- \mathcal{M} : the set of labeled canonical cotrees
- for any cotree t_0 with k leaves, \mathcal{M}_{t_0} : the set of labeled canonical cotrees with a marked k -tuple of distinct leaves, which induce t_0 .

with corresponding exponential generating series $M(z)$, $M_{t_0}(z)$,

Expressing $\mathbb{P}(\mathbf{t}_k^{(n)} = t_0)$

Notation:

- \mathcal{M} : the set of labeled canonical cotrees
- for any cotree t_0 with k leaves, \mathcal{M}_{t_0} : the set of labeled canonical cotrees with a marked k -tuple of distinct leaves, which induce t_0 .

with corresponding exponential generating series $M(z)$, $M_{t_0}(z)$,

$$\mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = \frac{n! [z^n] M_{t_0}(z)}{n! [z^n] M(z) \times n(n-1) \dots (n-k+1)}$$

Estimate the limit as $n \rightarrow \infty$ using analytic combinatorics,

Notation:

- \mathcal{M} : the set of labeled **canonical cotrees**
- for any cotree t_0 with k leaves, \mathcal{M}_{t_0} : the set of labeled canonical cotrees with a **marked** k -tuple of distinct **leaves**, which **induce** t_0 .
- \mathcal{L} : the set of non-plane rooted trees, labeled on their leaves, where internal nodes have ≥ 2 children.
Trees of \mathcal{L} are just like **cotrees without the decorations** on internal nodes.

with corresponding exponential generating series $M(z)$, $M_{t_0}(z)$, $L(z)$

$$\mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = \frac{n![z^n]M_{t_0}(z)}{n![z^n]M(z) \times n(n-1)\dots(n-k+1)}$$

Estimate the limit as $n \rightarrow \infty$ using **analytic combinatorics**,
on $L(z)$ and variants, **relating** $M(z)$ and $M_{t_0}(z)$ to $L(z)$

Behavior of $L(z)$ and $M(z)$

Study of $L(z)$:

From [Flajolet-Sedgewick] (rather a variant on trees counted by leaves):

- $L(z)$ satisfies $L(z) = z + \exp(L(z)) - 1 - L(z)$.

Behavior of $L(z)$ and $M(z)$

Study of $L(z)$:

From [Flajolet-Sedgewick] (rather a variant on trees counted by leaves):

- $L(z)$ satisfies $L(z) = z + \exp(L(z)) - 1 - L(z)$.
- $L(z)$ has radius of convergence $\rho = 2 \log(2) - 1$ and is Δ -analytic.
- Near $z = \rho$, $L(z) = \log(2) - \sqrt{\rho} \sqrt{1 - z/\rho} + \mathcal{O}(1 - z/\rho)$.

Behavior of $L(z)$ and $M(z)$

Study of $L(z)$:

From [Flajolet-Sedgewick] (rather a variant on trees counted by leaves):

- $L(z)$ satisfies $L(z) = z + \exp(L(z)) - 1 - L(z)$.
- $L(z)$ has radius of convergence $\rho = 2 \log(2) - 1$ and is Δ -analytic.
- Near $z = \rho$, $L(z) = \log(2) - \sqrt{\rho} \sqrt{1 - z/\rho} + \mathcal{O}(1 - z/\rho)$.

Study of $M(z)$:

- $M(z) = 2L(z) - z$, since all decorations in a canonical cotree (of size $\neq 1$) are determined by the decoration of the root (alternation).

Behavior of $L(z)$ and $M(z)$

Study of $L(z)$:

From [Flajolet-Sedgewick] (rather a variant on trees counted by leaves):

- $L(z)$ satisfies $L(z) = z + \exp(L(z)) - 1 - L(z)$.
- $L(z)$ has radius of convergence $\rho = 2 \log(2) - 1$ and is Δ -analytic.
- Near $z = \rho$, $L(z) = \log(2) - \sqrt{\rho} \sqrt{1 - z/\rho} + \mathcal{O}(1 - z/\rho)$.

Study of $M(z)$:

- $M(z) = 2L(z) - z$, since all decorations in a canonical cotree (of size $\neq 1$) are determined by the decoration of the root (alternation).
- $M(z)$ has radius of convergence $\rho = 2 \log(2) - 1$ and is Δ -analytic.
- Near $z = \rho$, $M(z) = 1 - 2\sqrt{\rho} \sqrt{1 - z/\rho} + \mathcal{O}(1 - z/\rho)$.

Behavior of $L(z)$ and $M(z)$

Study of $L(z)$:

From [Flajolet-Sedgewick] (rather a variant on trees counted by leaves):

- $L(z)$ satisfies $L(z) = z + \exp(L(z)) - 1 - L(z)$.
- $L(z)$ has radius of convergence $\rho = 2 \log(2) - 1$ and is Δ -analytic.
- Near $z = \rho$, $L(z) = \log(2) - \sqrt{\rho} \sqrt{1 - z/\rho} + \mathcal{O}(1 - z/\rho)$.

Study of $M(z)$:

- $M(z) = 2L(z) - z$, since all decorations in a canonical cotree (of size $\neq 1$) are determined by the decoration of the root (alternation).
- $M(z)$ has radius of convergence $\rho = 2 \log(2) - 1$ and is Δ -analytic.
- Near $z = \rho$, $M(z) = 1 - 2\sqrt{\rho} \sqrt{1 - z/\rho} + \mathcal{O}(1 - z/\rho)$.
- From the [transfer theorem](#),

$$n(n-1) \dots (n-k+1) [z^n] M(z) \underset{n \rightarrow +\infty}{\sim} \frac{n^{k-3/2}}{\rho^{n-1/2} \sqrt{\pi}}.$$

Recall: Trees of \mathcal{M}_{t_0} are trees of \mathcal{L} with k marked leaves inducing t_0 , and in addition a decoration on the root

Recall: Trees of \mathcal{M}_{t_0} are trees of \mathcal{L} with k marked leaves inducing t_0 , and in addition a decoration on the root

Terminology:

- Marked leaf: carries a label (hence contributes 1 to the size)
- Blossom: similar to a marked leaf, but does not carry a label

Recall: Trees of \mathcal{M}_{t_0} are trees of \mathcal{L} with k marked leaves inducing t_0 , and in addition a decoration on the root

Terminology:

- Marked leaf: carries a label (hence contributes 1 to the size)
- Blossom: similar to a marked leaf, but does not carry a label

Series to consider:

- $L'(z)$: counts trees of \mathcal{L} with a blossom
- $L^\bullet(z) = zL'(z)$: counts trees of \mathcal{L} with a marked leaf
- L^{odd} : counts trees of \mathcal{L} with a blossom at odd distance from the root
- L^{even} : same for even distance

Recall: Trees of \mathcal{M}_{t_0} are trees of \mathcal{L} with k marked leaves inducing t_0 , and in addition a decoration on the root

Terminology:

- Marked leaf: carries a label (hence contributes 1 to the size)
- Blossom: similar to a marked leaf, but does not carry a label

Series to consider:

- $L'(z)$: counts trees of \mathcal{L} with a blossom
- $L^\bullet(z) = zL'(z)$: counts trees of \mathcal{L} with a marked leaf
- L^{odd} : counts trees of \mathcal{L} with a blossom at odd distance from the root
- L^{even} : same for even distance

Simple combinatorial arguments give $L^{\text{even}} = \frac{1}{e^L(2-e^L)}$ and $L^{\text{odd}} = \frac{e^L-1}{e^L(2-e^L)}$.

Recall: Trees of \mathcal{M}_{t_0} are trees of \mathcal{L} with k marked leaves inducing t_0 , and in addition a decoration on the root

Terminology:

- Marked leaf: carries a label (hence contributes 1 to the size)
- Blossom: similar to a marked leaf, but does not carry a label

Series to consider:

- $L'(z)$: counts trees of \mathcal{L} with a blossom
- $L^\bullet(z) = zL'(z)$: counts trees of \mathcal{L} with a marked leaf
- L^{odd} : counts trees of \mathcal{L} with a blossom at odd distance from the root
- L^{even} : same for even distance

Simple combinatorial arguments give $L^{\text{even}} = \frac{1}{e^{L(2-e^L)}}$ and $L^{\text{odd}} = \frac{e^L - 1}{e^{L(2-e^L)}}$.

Thus, the singular behavior of $L(z)$ determines the one of these four series.


Prop.: If t_0 with k leaves has n_v internal vertices, $n_=\$ edges of the form $0 - 0$ or $1 - 1$, and n_\neq edges of the form $0 - 1$ or $1 - 0$, then

$$M_{t_0} = (L')(\exp(L))^{n_v}(L^\bullet)^k(L^{\text{odd}})^{n_=(L^{\text{even}})^{n_\neq}}.$$

Prop.: If t_0 with k leaves has n_v internal vertices, $n_=\$ edges of the form $0 - 0$ or $1 - 1$, and n_{\neq} edges of the form $0 - 1$ or $1 - 0$, then

$$M_{t_0} = (L')(\exp(L))^{n_v} (L^\bullet)^k (L^{\text{odd}})^{n_=} (L^{\text{even}})^{n_{\neq}}.$$

Proof:



Prop.: If t_0 with k leaves has n_v internal vertices, $n_=\$ edges of the form $0 - 0$ or $1 - 1$, and n_{\neq} edges of the form $0 - 1$ or $1 - 0$, then

$$M_{t_0} = (L')(\exp(L))^{n_v} (L^\bullet)^k (L^{\text{odd}})^{n_=} (L^{\text{even}})^{n_{\neq}}.$$

Proof:



Corollary: Like before, we obtain

- the behavior at ρ of $M_{t_0}(z)$,
- and the asymptotic estimate of $[z^n]M_{t_0}(z)$.

Prop.: If t_0 with k leaves has n_v internal vertices, $n_ =$ edges of the form $0 - 0$ or $1 - 1$, and n_{\neq} edges of the form $0 - 1$ or $1 - 0$, then

$$M_{t_0} = (L')(\exp(L))^{n_v} (L^\bullet)^k (L^{\text{odd}})^{n_ =} (L^{\text{even}})^{n_{\neq}}.$$

Proof:



Corollary: Like before, we obtain

- the behavior at ρ of $M_{t_0}(z)$,
- and the asymptotic estimate of $[z^n]M_{t_0}(z)$.

More precisely, we have

$$[z^n]M_{t_0}(z) \underset{n \rightarrow +\infty}{\sim} \frac{(k-1)!}{(2k-2)!} \frac{n^{k-3/2}}{\rho^{n-1/2} \sqrt{\pi}},$$

if t_0 is binary (which implies $n_v = k - 1$ and $n_ + n_{\neq} = k - 2$).

Conclusion of the combinatorial study (labeled case)

Notation (reminder):

- $\mathbf{t}^{(n)}$: uniform random labeled canonical cotree of size n
- $\mathbf{t}_k^{(n)}$: subtree of $\mathbf{t}^{(n)}$ induced by a uniform k -tuple of distinct leaves
- t_0 : cotree with k leaves

What we proved: If t_0 is **binary**, then $\lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = \frac{(k-1)!}{(2k-2)!}$.

Conclusion of the combinatorial study (labeled case)

Notation (reminder):

- $\mathbf{t}^{(n)}$: uniform random labeled canonical cotree of size n
- $\mathbf{t}_k^{(n)}$: subtree of $\mathbf{t}^{(n)}$ induced by a uniform k -tuple of distinct leaves
- t_0 : cotree with k leaves

What we proved: If t_0 is **binary**, then $\lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = \frac{(k-1)!}{(2k-2)!}$.

Remark: $\frac{(k-1)!}{(2k-2)!} = \frac{1}{\text{number of binary cotrees with } k \text{ leaves}}$.

Consequence: If t_0 is **not binary**, then $\lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = 0$.

Conclusion of the combinatorial study (labeled case)

Notation (reminder):

- $\mathbf{t}^{(n)}$: uniform random labeled canonical cotree of size n
- $\mathbf{t}_k^{(n)}$: subtree of $\mathbf{t}^{(n)}$ induced by a uniform k -tuple of distinct leaves
- t_0 : cotree with k leaves

What we proved: If t_0 is **binary**, then $\lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = \frac{(k-1)!}{(2k-2)!}$.

Remark: $\frac{(k-1)!}{(2k-2)!} = \frac{1}{\text{number of binary cotrees with } k \text{ leaves}}$.

Consequence: If t_0 is **not binary**, then $\lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{t}_k^{(n)} = t_0) = 0$.

Remark/reminder:

Summing over all t_0 encoding a cograph g , this gives $\lim_{n \rightarrow \infty} \mathbb{E}[\text{Dens}(g, \mathbf{G}_n)]$.


**The Brownian cographon
and its expected subgraph densities
(or something close to it)**

Defining the Brownian cographon

Decorated Brownian excursion:

- \mathbf{e} : Brownian excursion of length 1.
- $(\mathbf{b}_i)_{i \geq 1}$: enumeration of the locations of the local minima of \mathbf{e} (which exists).
- $\mathbf{S}^p = (\mathbf{s}_1, \dots)$: sequence of i.i.d. r.v. in $\{0, 1\}$, independent from \mathbf{e} , with $\mathbb{P}(\mathbf{s}_1 = 0) = p$.


In the decorated Brownian excursion $(\mathbf{e}, \mathbf{S}^p)$, we think of the decorations \mathbf{s}_j as attached to the local minimum at \mathbf{b}_j .



Defining the Brownian cographon

Decorated Brownian excursion:

- \mathbf{e} : Brownian excursion of length 1.
- $(\mathbf{b}_i)_{i \geq 1}$: enumeration of the locations of the local minima of \mathbf{e} (which exists).
- $\mathbf{S}^p = (\mathbf{s}_1, \dots)$: sequence of i.i.d. r.v. in $\{0, 1\}$, independent from \mathbf{e} , with $\mathbb{P}(\mathbf{s}_1 = 0) = p$.




In the decorated Brownian excursion $(\mathbf{e}, \mathbf{S}^p)$, we think of the decorations \mathbf{s}_i as attached to the local minimum at \mathbf{b}_i .

Brownian cographon of parameter $p \in [0, 1]$, \mathbf{W}^p :

- for any $x, y \in [0, 1]$, $\text{Dec}(x, y; \mathbf{e}, \mathbf{S}^p) \in \{0, 1\} =$ decoration of the local minimum of \mathbf{e} on $[x, y]$ (or $[y, x]$) (a.s. unique and $\neq x, y$)
- $\mathbf{W}^p =$ graphon associated with the function
$$\begin{aligned} \mathbf{w}^p : [0, 1]^2 &\rightarrow \{0, 1\}; \\ (x, y) &\mapsto \text{Dec}(x, y; \mathbf{e}, \mathbf{S}^p). \end{aligned}$$

“Here is” $W^{1/2}$



This is actually the adjacency matrix of a uniform random labeled cograph of size 4482, where the order of the vertices to plot the matrix is the depth-first search on the associated cotree.

Distribution of induced subgraphs of W^p

Notation:

- W^p : Brownian cographon of parameter p
- $\text{Sample}_k(W)$: subgraph of W induced by k i.i.d. uniform “vertices” $x_1, \dots, x_k \in [0, 1]$
- \mathbf{b}_k^p : uniform labeled binary tree with k leaves, where internal vertices carry $\{0, 1\}$ decorations with $\mathbb{P}(0) = p$.

Prop.: $\text{Sample}_k(W^p) \stackrel{(d)}{=} \text{Cograph}(\mathbf{b}_k^p)$.

Distribution of induced subgraphs of W^p


Notation:

- W^p : Brownian cographon of parameter p
- $\text{Sample}_k(W)$: subgraph of W induced by k i.i.d. uniform “vertices” $x_1, \dots, x_k \in [0, 1]$
- \mathbf{b}_k^p : uniform labeled binary tree with k leaves, where internal vertices carry $\{0, 1\}$ decorations with $\mathbb{P}(0) = p$.

Prop.: $\text{Sample}_k(W^p) \stackrel{(d)}{=} \text{the unlabeled version of } \text{Cograph}(\mathbf{b}_k^p)$.

Proof idea:

- \mathbf{b}_k^p is the cotree extracted from $(\mathbf{e}, \mathbf{S}^p)$ and x_1, \dots, x_k .
- $\text{Sample}_k(W^p)$ is the associated cograph since decorations indicate edges similarly in W^p and in $\text{Cograph}(\mathbf{b}_k^p)$.



Characterization of convergence to $\mathcal{W}^{1/2}$

Prop.: For $(\mathbf{t}^{(n)})_n$ a sequence of random cotrees s.t. $\text{size}(\mathbf{t}^{(n)}) = n$,
let $\mathbf{t}_k^{(n)}$ be the subtree of $\mathbf{t}^{(n)}$ induced by a unif. k -tuple of distinct leaves.
If for any binary cotree t_0 we have $\mathbb{P}(\mathbf{t}_k^{(n)} = t_0) \xrightarrow{n \rightarrow \infty} \frac{(k-1)!}{(2k-2)!}$, (★)
then $(\text{Cograph}(\mathbf{t}^{(n)}))_n$ converges to $\mathcal{W}^{1/2}$.

Characterization of convergence to $\mathcal{W}^{1/2}$

Prop.: For $(\mathbf{t}^{(n)})_n$ a sequence of random cotrees s.t. $\text{size}(\mathbf{t}^{(n)}) = n$, let $\mathbf{t}_k^{(n)}$ be the subtree of $\mathbf{t}^{(n)}$ induced by a unif. k -tuple of distinct leaves.

If for any binary cotree t_0 we have $\mathbb{P}(\mathbf{t}_k^{(n)} = t_0) \xrightarrow{n \rightarrow \infty} \frac{(k-1)!}{(2k-2)!}$, (★)

then $(\text{Cograph}(\mathbf{t}^{(n)}))_n$ converges to $\mathcal{W}^{1/2}$.

Proof idea:

- (★) says $\mathbf{t}_k^{(n)}$ is asymptotically uniform on labeled binary cotrees with k leaves, which is distributed like $\mathbf{b}_k^{1/2}$
- Take **cographs** and forget labels
 $\Rightarrow \text{SubGraph}_k^{\text{inj}}(\text{Cograph}(\mathbf{t}^{(n)})) \xrightarrow{(d)} \text{Sample}_k(\mathcal{W}^{1/2})$

Characterization of convergence to $\mathcal{W}^{1/2}$

Prop.: For $(\mathbf{t}^{(n)})_n$ a sequence of random cotrees s.t. $\text{size}(\mathbf{t}^{(n)}) = n$, let $\mathbf{t}_k^{(n)}$ be the subtree of $\mathbf{t}^{(n)}$ induced by a unif. k -tuple of distinct leaves. If for any binary cotree t_0 we have $\mathbb{P}(\mathbf{t}_k^{(n)} = t_0) \xrightarrow{n \rightarrow \infty} \frac{(k-1)!}{(2k-2)!}$, (\star) then $(\text{Cograph}(\mathbf{t}^{(n)}))_n$ converges to $\mathcal{W}^{1/2}$.

Proof idea:

- (\star) says $\mathbf{t}_k^{(n)}$ is asymptotically uniform on labeled binary cotrees with k leaves, which is distributed like $\mathbf{b}_k^{1/2}$
- Take **cographs** and forget labels
 $\Rightarrow \text{SubGraph}_k^{\text{inj}}(\text{Cograph}(\mathbf{t}^{(n)})) \xrightarrow{(d)} \text{Sample}_k(\mathcal{W}^{1/2})$

Corollary: (\mathbf{G}_n) converges to $\mathcal{W}^{1/2}$.

Uniform random labeled cographs converge to the Brownian cographon.

(apply the prop. to $\mathbf{t}^{(n)} =$ unif. random canonical labeled cotree of size n)

Additional results

Degree distribution of graphs and graphons:

- (Rescaled) degree distribution of G : $D_G = \frac{1}{n} \sum_{v \text{ vertex}} \delta_{\deg(v)/n}$
- It generalizes to graphons: for w representing W , D_W is defined by $\int_{[0,1]} f(x) D_W(dx) = \int_{[0,1]} f \left(\int_{[0,1]} w(u, v) dv \right) du, \forall f \text{ cont. bounded}$
- D_G and D_W are probability measures on $[0, 1]$

Lemma: If $(G_n)_n$ converges to W , then (D_{G_n}) converges (weakly) to D_W .

Degree distribution of graphs and graphons:

- (Rescaled) degree distribution of G : $D_G = \frac{1}{n} \sum_{v \text{ vertex}} \delta_{\deg(v)/n}$
- It generalizes to graphons: for w representing W , D_W is defined by $\int_{[0,1]} f(x) D_W(dx) = \int_{[0,1]} f \left(\int_{[0,1]} w(u, v) dv \right) du, \forall f \text{ cont. bounded}$
- D_G and D_W are probability measures on $[0, 1]$

Lemma: If $(G_n)_n$ converges to W , then (D_{G_n}) converges (weakly) to D_W .

With W random: D_W is a random measure.

Its intensity measure $I[D_W]$ is the “averaged” degree distribution of W , where we average on all realizations of W

Degree distribution of graphs and graphons:

- (Rescaled) degree distribution of G : $D_G = \frac{1}{n} \sum_{v \text{ vertex}} \delta_{\deg(v)/n}$
- It generalizes to graphons: for w representing W , D_W is defined by $\int_{[0,1]} f(x) D_W(dx) = \int_{[0,1]} f \left(\int_{[0,1]} w(u, v) dv \right) du, \forall f \text{ cont. bounded}$
- D_G and D_W are probability measures on $[0, 1]$

Lemma: If $(G_n)_n$ converges to W , then (D_{G_n}) converges (weakly) to D_W .

With W random: D_W is a random measure.

Its intensity measure $I[D_W]$ is the “averaged” degree distribution of W , where we average on all realizations of W

Prop.: For the Brownian cographon, $I[D_{W_{1/2}}]$ is uniform on $[0, 1]$.

Corollary: The rescaled degree of a uniform random vertex v_n in G_n is asymptotically uniform in $[0, 1]$.

Same results:

- *Definition:* \mathbf{G}_n^u = uniform random **unlabeled** cograph with n vertices
- *Theorem:* $(\mathbf{G}_n^u)_n$ **converges to** the Brownian cographon $\mathbf{W}^{1/2}$
- *Consequence:* The **rescaled degree** of a uniform random vertex \mathbf{v}_n in \mathbf{G}_n^u is **asymptotically uniform** in $[0, 1]$.

The unlabeled case

Same results:

- *Definition:* \mathbf{G}_n^u = uniform random **unlabeled** cograph with n vertices
- *Theorem:* $(\mathbf{G}_n^u)_n$ **converges to** the Brownian cographon $\mathbf{W}^{1/2}$
- *Consequence:* The **rescaled degree** of a uniform random vertex \mathbf{v}_n in \mathbf{G}_n^u is **asymptotically uniform** in $[0, 1]$.

How to modify the proof:

- Same strategy of analytic combinatorics, using **unlabeled** cotrees.
- With **Pólya operators**, it is difficult to count objects **with marked leaves** (inducing a given subtree t_0).
- Instead of \mathcal{L} as before, we study

$$\mathcal{U} = \{(t, a) : t \in \mathcal{L}, a \text{ a root-preserving automorphism of } t\}.$$

Using \mathcal{U} , we can interpret Pólya operators **combinatorially**, in a way that allows to **keep track of marked leaves**.

Remark: For their **graphon limit** (and average degree distribution), labeled and unlabeled cographs display the **same** behavior.

Question: Are there some statistics which behave **differently** in the labeled and unlabeled case?

Remark: For their **graphon limit** (and average degree distribution), labeled and unlabeled cographs display the **same** behavior.

Question: Are there some statistics which behave **differently** in the labeled and unlabeled case?

Example of the vertex connectivity:

- $\kappa(G)$ = minimal number of vertices whose removal disconnects G
- For a connected cograph G with canonical cotree T (with root 1), $\kappa(G) = |G| - |T_{max}|$, where $|T_{max}|$ is the largest component of T
- Using again analytic combinatorics, we express, for all $j \geq 1$,
 $\lim_{n \rightarrow \infty} \mathbb{P}(\kappa(\mathbf{G}_n) = j)$ using $L(z)$ as $1/2 \cdot \rho_L^j[z^j] (e^{L(z)} - 1)$
 $\lim_{n \rightarrow \infty} \mathbb{P}(\kappa(\mathbf{G}_n^u) = j)$ using $U(z)$ as $1/2 \cdot \rho_U^j[z^j] (2U(z) - z)$
(the limiting probability of having $\kappa(\mathbf{G}_n)$ or $\kappa(\mathbf{G}_n^u) = 0$ being $1/2$).
- These limit distributions are **different**.

A parallel with permutations *via* inversion graphs

Separable permutations

- encoding by decomposition trees
- convergence to the Brownian separable permuton (BSP)
- [BBFGP18, Maazoun16, BBFS20]

Cographs

- encoding by cotrees
- convergence to the Brownian Cographon (BCG)
- [this talk, Stufler21]

A parallel with permutations *via* inversion graphs

Separable permutations

- encoding by decomposition trees
- convergence to the Brownian separable permuton (BSP)
- [BBFGP18, Maazoun16, BBFS20]

Substitution-closed classes

- encoding by decomposition trees
- Universality of the BSP
- [BBFGMP20]

Beyond these families

- encoding by decomposition trees
- BSP VS X-permuton
- [BBFGMP21+]

Cographs

- encoding by cotrees
- convergence to the Brownian Cographon (BCG)
- [this talk, Stufler21]

A parallel with permutations *via* inversion graphs

Separable permutations

- encoding by decomposition trees
- convergence to the Brownian separable permutation (BSP)
- [BBFGP18, Maazoun16, BBFS20]

Substitution-closed classes

- encoding by decomposition trees
- Universality of the BSP
- [BBFGMP20]

Beyond these families

- encoding by decomposition trees
- BSP VS X-permutation
- [BBFGMP21+]

Cographs

- encoding by cotrees
- convergence to the Brownian Cographon (BCG)
- [this talk, Stufler21]

Classes of graphs closed for the substitution operation of the modular decomposition

- encoding by modular decomposition trees
- Expected universality of the BCG for “small” classes

Beyond these families

- Other graph decompositions?

(Spoiler: The next result of the (extended) team [BBDFGMP])

Results:

- The size of the **largest independent set** of a uniform random cograph is **sublinear**.
(hence P_4 does not have the asymptotic linear Erdős-Hajnal property)
- The length of the **longest increasing subsequence** of a uniform random separable permutations is **sublinear**.

Independence number and longest increasing subsequences

(Spoiler: The next result of the (extended) team [BBDFGMP])

Results:

- The size of the **largest independent set** of a uniform random cograph is **sublinear**.
(hence P_4 does not have the asymptotic linear Erdős-Hajnal property)
- The length of the **longest increasing subsequence** of a uniform random separable permutations is **sublinear**.

Main proof ingredients:

- Convergence to the **Brownian cographon**
- The independence number of the Brownian cographon $\mathbf{W}^{1/2}$ is 0

Independence number and longest increasing subsequences

(Spoiler: The next result of the (extended) team [BBDFGMP])

Results:

- The size of the **largest independent set** of a uniform random cograph is **sublinear**.
(hence P_4 does not have the asymptotic linear Erdős-Hajnal property)
- The length of the **longest increasing subsequence** of a uniform random separable permutations is **sublinear**.

Main proof ingredients:

- Convergence to the **Brownian cographon**
- The independence number of the Brownian cographon $\mathbf{W}^{1/2}$ is 0

Bonus: The sublinearity result applies to **all classes** with graphon/permuton limit \mathbf{W}^P or a Brownian separable permuton.

Independence number and longest increasing subsequences

(Spoiler: The next result of the (extended) team [BBDFGMP])

Results:

- The size of the **largest independent set** of a uniform random cograph is **sublinear**.
(hence P_4 does not have the asymptotic linear Erdős-Hajnal property)
- The length of the **longest increasing subsequence** of a uniform random separable permutations is **sublinear**.

Main proof ingredients:

- Convergence to the **Brownian cographon**
- The independence number of the Brownian cographon $\mathbf{W}^{1/2}$ is 0

Bonus: The sublinearity result applies to **all classes** with graphon/permuton limit \mathbf{W}^P or a Brownian separable permuton.

Thank you for being there!